

The Journal of the Royal Society for the Promotion of Health

<http://rsh.sagepub.com>

Central obesity and coronary risk factors

Arnab Ghosh, Kaushik Bose, Shila Chakravarti, Asit Baran Das Chaudhuri, Jayanti Chattopadhyay, Gautam Dasgupta and Sivaji Sengupta

The Journal of the Royal Society for the Promotion of Health 2004; 124; 86

DOI: 10.1177/146642400412400213

The online version of this article can be found at:
<http://rsh.sagepub.com/cgi/content/abstract/124/2/86>

Published by:



<http://www.sagepublications.com>

On behalf of:



[Royal Society for the Promotion of Health](http://www.rsh.sagepub.com)

Additional services and information for *The Journal of the Royal Society for the Promotion of Health* can be found at:

Email Alerts: <http://rsh.sagepub.com/cgi/alerts>

Subscriptions: <http://rsh.sagepub.com/subscriptions>

Reprints: <http://www.sagepub.com/journalsReprints.nav>

Permissions: <http://www.sagepub.com/journalsPermissions.nav>

Citations (this article cites 24 articles hosted on the SAGE Journals Online and HighWire Press platforms):
<http://rsh.sagepub.com/cgi/content/refs/124/2/86>

Central obesity and coronary risk factors

Authors

Amab Ghosh, Department of Anthropology, University of Calcutta, Kolkata, India

Kaushik Bose, Department of Anthropology, Vidyasagar University, Midnapore, India

Shila Chakravarti, Department of Anthropology, University of Calcutta, as above

Asit Baran Das Chaudhuri, Department of Anthropology, University of Calcutta, as above

Jayanti Chattopadhyay, Department of Pathology, BR Singh Hospital, Kolkata, India

Gautam Dasgupta, Department of Pathology, BR Singh Hospital, as above

Sivaji Sengupta, Department of Pathology, BR Singh Hospital, as above

Corresponding author:

Kaushik Bose, Reader in Biological Anthropology, Department of Anthropology, Vidyasagar University Midnapore 721 102, West Bengal, India Email: banda@vsnl.net

Full title: Relationship of waist circumference and waist-hip ratio with metabolic risk factors of coronary heart disease among Bengalee Hindu men of Kolkata, India

Received 20 August 2001, revised and accepted 11 June 2002

Key words

Bengalees; central obesity; metabolic variables

Abstract

A cross-sectional study of 130 Bengalee Hindu men (mean age=50.3 years; SD=10.5 years) was undertaken to investigate the relationship of body mass index (BMI), waist circumference (WC) and waist-hip ratio (WHR) with total cholesterol (TC), high density (HDL-C), low density (LDL-C) and very low-density (VLDL-C) lipoprotein cholesterol, fasting plasma glucose (FPG) and triglycerides (FTG). Correlation studies revealed that WHR was significantly correlated ($r=0.245$, $p<0.01$) with TC. WC and WHR had significant correlations with VLDL-C, FPG and FTG. All subjects were further divided into two groups based on $WHR \leq 0.95$ (centrally non-obese, CNO) and $WHR > 0.95$ (centrally obese, CO) following the US Joint National Committee (JNC) guidelines. Students' t-test revealed that CO subjects ($n=83$) had a significantly higher mean TC ($p<0.05$), VLDL ($p<0.05$), FPG ($p<0.01$) and FTG ($p<0.05$) compared with CNO individuals ($n=47$). Results of analysis of variance (ANOVA) of central obesity status (CNO=no, CO=yes) and BMI (BMI tertiles used as a categorical variable) with these metabolic variables revealed that CO status had a significant effect ($p<0.05$) on TC, VLDL-C, FPG and FTG. BMI tertiles did not have significant effect on any of these metabolic variables. There was no significant BMI tertile-central obesity status interaction. It can therefore be concluded that the JNC guidelines of $WHR > 0.95$ to define central obesity can be used, irrespective of BMI, among this population, to identify individuals who have enhanced metabolic risk factors of coronary heart disease (CHD). Furthermore, it can be routinely used for health promotion purposes among Bengalee men.

INTRODUCTION

Obesity is now so common within the world's population that it is beginning to replace undernutrition and infectious diseases as the most significant contributor to ill health. In particular, obesity is associated with type 2 diabetes, coronary heart disease (CHD) and other associated disorders.¹ Vague² first suggested that body fat around the abdomen was related to diabetes and atherosclerosis in both men and women. However, there is no universally agreed way of measuring adiposity, nor is it known which measure is the best predictor of cardiovascular disease.

Several researchers have favoured waist to hip ratio (WHR) arguing that it is a simple, useful and better indicator of risk factors and a better predictor of CHD³⁻⁶ than other used measures of adiposity such as body mass index (BMI) and waist to height ratio. In some recent studies⁷ waist circumference (WC) has also been utilised to measure central adiposity. The relationship of WHR with risk factors of CHD has been investigated in different ethnic groups: native Americans,^{8,9} Mexican-Americans,^{7,10} African-Americans,¹¹ Melanesians,¹²

and Central Asians.¹³ These studies show that the association of central adiposity with metabolic risk factors is not the same at all levels of adiposity in all ethnic groups. A recent report of the Joint National Committee (JNC)¹⁴ has suggested that a $WHR > 0.95$ in males is associated with a higher coronary risk.

Asian Indians, both in India as well as migrants elsewhere, have unusually high rates of CHD.^{15,16} Studies have investigated the relationship of BMI¹⁷ and WHR¹⁸⁻²⁰ with risk factors of CHD among migrant South Asians. Although recent studies from India have investigated the association of WHR with hypertension²¹ and other risk factors for CHD,^{22,23} none of these studies have used the JNC cut-off point of $WHR > 0.95$ to classify individuals as centrally obese. Although Dasgupta and Hazra²⁴ utilised the JNC cut-off point of $WHR > 0.95$ to determine the prevalence of central obesity among Bengalee men and women, they did not study the relation of central obesity with metabolic risk factors of CHD. Therefore, no information is available on the relationship of WC and WHR with metabolic risk factors of CHD among Bengalees. The present study on Bengalee Hindu men had two

objectives: firstly, to investigate the relationship of BMI, WC and WHR with some metabolic risk factors of CHD; and secondly, to investigate the relationship of central obesity, at different levels of BMI, on metabolic risk of CHD.

METHODS

Study population

The present study was conducted during December 1999–July 2000 at the Outpatients Department of BR Singh Hospital, Eastern Railways, Calcutta, as part of a collaborative research programme between the Department of Anthropology, University of Calcutta and the Department of Pathology, BR Singh Hospital. Prior to the commencement of the study, written information regarding the aims, objectives and the criteria for eligibility were sent to all employees of the Eastern Railways at Kolkata. Any employee who belonged to the Hindu caste population and was more than 30 years of age was eligible to participate in this health check-up programme. The study sample was therefore representative of the general Bengalee Hindu male population.

Anthropometric and lipid profile measurements were made after the subjects had completed a questionnaire which requested information on their age, occupation, medical history, exercise undertaken and alcohol consumption. Only two subjects were receiving treatment either for high cholesterol or diabetes and they were excluded from the analyses. A total of 14 men had undiagnosed diabetes and they were included in the study since they formed a representative part of the population. Their inclusion did not bias the results when the data were re-analysed without them. The sample size of this study was 130.

Anthropometric measurements

Height, weight, waist and hip circumference measurements were made using standard techniques of Lohman *et al*²⁵ by a trained investigator (AG). Height and weight were measured to the nearest 0.1 cm and 0.5 kg, respectively. Waist and hip circumferences were measured with a tape to the nearest 0.2 cm. BMI and WHR were computed using the standard formulae:

$\text{WHR} = \text{waist circumference (cm)} / \text{hip}$

$\text{circumference (cm)}$

$\text{BMI} = \text{weight (kg)} / \text{height}^2 \text{ (m}^2\text{)}$

The JNC¹⁴ criterion of $\text{WHR} > 0.95$ was utilised as a cut-off point to define central obesity as suggested by several researchers.^{24, 26} It is now widely accepted²⁶ that a $\text{WHR} > 0.95$ in men is associated with lipid abnormalities and confers the highest risk of CHD. The JNC guidelines classify men into two groups: centrally non-obese (CNO) and centrally obese (CO). The cut-off points used were: $\text{CNO} = \text{WHR} \leq 0.95$; $\text{CO} = \text{WHR} > 0.95$. There were 47 and 83 subjects, respectively, in these two groups.

Metabolic variables

A fasting blood sample was collected from each subject for the determination of

metabolic variables. All subjects maintained an overnight fast (at least 12 hours duration) prior to blood collection. Plasma was separated by centrifugation at $1,000 \times g$ for 20 min at room temperature within two hours of collection. Estimation of total cholesterol (TC), fasting plasma glucose (FPG) and fasting triglyceride (FTG) were carried out on separated plasma using a Technicon RA-XT autoanalyzer (Technicon Instruments Corporation, NY, USA). High-density lipoprotein cholesterol (HDL-C) was measured after an overnight stand of plasma in a refrigerator and then precipitation of non-high-density lipoproteins (LDL, VLDL, chylomicrons) with manganese-heparin substrate.²⁷

Values of low-density lipoprotein cholesterol (LDL-C) and very low-density lipoprotein cholesterol (VLDL-C) were

Table 1

Anthropometric and metabolic characteristics of the study population

Variable	Mean	SD
Age (years)	50.3	10.5
Anthropometric		
Height (cm)	164.3	7.12
Weight (kg)	64.3	9.17
BMI (kg/m ²)	23.8	2.80
WC (cm)	86.0	7.00
HC (cm)	89.7	5.14
WHR	0.959	0.046
Metabolic		
TC (mmol/l)	5.5	0.92
HDL-C (mmol/l)	1.2	0.09
LDL-C (mmol/l)	3.3	0.74
VLDL-C (mmol/l)*	0.3	0.22
HDL/LDL ratio	0.4	0.09
FPG (mmol/l)	6.2	1.10
FTG (mmol/l)*	1.9	0.22

*Geometric means are presented

SD = standard deviation

HC = hip circumference

TC = total cholesterol

FPG = fasting blood glucose

WC = waist circumference

VLDL-C = very low density lipoprotein cholesterol

HDL-C = high density lipoprotein cholesterol

LDL-C = low density lipoprotein cholesterol

WHR = waist-hip ratio

FTG = fasting triglyceride

estimated using the following formulae²⁸:

$$\text{LDL-C} = \text{TC} - (\text{HDL-C} + \text{FTG}/5)$$

$$\text{VLDL-C} = \text{FTG}/5$$

All biochemical analyses were carried out at the Biochemistry Unit of the Department of Pathology, BR Singh Hospital, Kolkata. All metabolic variables were measured in mg/dl and then converted into mmol/l by using the following standard conversion formula:

For TC, HDL-C, LDL-C and VLDL-C:
value in mg/dl \times 0.02586

For FTG: value in mg/dl \times 0.01129

For FPG: value in mg/dl \times 0.05551

Statistical analyses

The distributions of all variables and indices were checked for normality. The distributions of VLDL-C and FTG were significantly skewed. Log(10) transformations were undertaken to normalise their distributions. All statistical analyses of these two variables were performed on log(10) transformed values. Correlation studies were undertaken to investigate the associations of BMI, WC, WHR and the metabolic variables. Metabolic differences between CNO and CO were studied using Student's t-test. Finally, analysis of variance (ANOVA) was utilised to test the impact of BMI (BMI tertiles were used, first tertile=22.7 kg/m², second tertile=24.8 kg/m²) and central obesity

status (categorical variable: CNO=no; CO=yes) on metabolic variables. All statistical analyses were performed using the SPSS (Statistical Package for Social Sciences) Version 5 Package. Statistical significance was set at $p < 0.05$.

RESULTS

The mean age of the subjects was 50.3 years (SD=10.5 years). The majority of the subjects (96%) had a sedentary lifestyle. None of them consumed alcohol. There was no significant relationship between age and the anthropometric variables and indices. Table 1 presents the means and standard deviations of the anthropometric and metabolic variables and indices of the subjects. Geometric means of VLDL-C and FTG are presented.

The associations of BMI, WC, WHR and the metabolic variables were studied utilising correlation analyses. Results (Table 2) revealed that WHR was significantly associated with TC ($r=0.245$), VLDL-C ($r=0.230$), FPG ($r=0.177$) and FTG ($r=0.198$). WC was significantly associated with VLDL-C ($r=0.174$), FPG ($r=0.177$) and FTG ($r=0.175$). BMI did not have significant association with any of the metabolic variables.

Since it was observed that WHR had significant relationship with various metabolic variables (Table 2), subjects were divided into two groups (CNO, $n=47$ and CO, $n=83$) following the JNC¹⁴ criterion to compare their metabolic profiles. The mean age of both groups was similar (CNO=50.5 years, SD=9.6; CO=50.2 years, SD=11.1). Students' t-test revealed that

CO subjects had significantly higher mean TC ($p < 0.05$), VLDL-C ($p < 0.05$), FPG ($p < 0.01$) and FTG ($p < 0.05$) compared with CNO individuals.

Since there were significant differences between CNO and CO subjects in TC, VLDL, FPG and FTG, ANOVA was undertaken to study the effect of BMI (BMI tertiles used as a categorical variable) and central obesity status (CNO=no; CO=yes) on these metabolic variables. Results (Table 3) revealed that CO status had significant effect ($p < 0.05$) on TC ($F=4.270$), VLDL ($F=5.488$), FPG ($F=5.826$) and FTG ($F=5.488$). BMI did not have significant effect on any of these variables. Furthermore, there was no significant BMI-central obesity interaction for any of these metabolic variables. Moreover, CO subjects had greater mean TC, VLDL-C, FPG and FTG compared to the overall means. In each case, CNO individuals had lower means compared to the overall means.

DISCUSSION

Obesity causes or exacerbates many health problems, both independently and in association with other diseases.²⁹ Obese individuals with excess intra-abdominal fat are at particular risk of negative health consequences, with certain ethnic populations like migrant Indians carrying different levels of risk.¹⁸ Total body fat appears to be a less important indicator of metabolic complications than the fat distribution pattern.

Although there is no universally agreed way of measuring central adiposity, WHR

Table 2

Pearson correlation coefficients (r) of BMI, WC and WHR with metabolic variables

	BMI		WC		WHR	
	r	p	r	p	r	p
TC	0.005	0.954	0.117	0.185	0.245	0.005
HDL-C	-0.075	0.398	-0.001	0.995	0.115	0.193
LDL-C	0.008	0.926	0.010	0.909	0.137	0.119
VLDL-C	0.051	0.565	0.174	0.048	0.230	0.020
FPG	0.149	0.091	0.177	0.042	0.177	0.042
FTG	0.047	0.599	0.175	0.046	0.198	0.024

BMI = body mass index

WC = waist circumference

WHR = waist-hip ratio

TC = total cholesterol

HDL-C = high density lipoprotein cholesterol

LDL-C = low density lipoprotein cholesterol

VLDL-C = very low density lipoprotein cholesterol

FPG = fasting blood glucose

FTG = fasting triglyceride

has been found to be the best predictor of cardiovascular disease.^{5,6,26} A recent report¹⁴ has suggested that a WHR>0.95 in males and WHR>0.85 in females is associated with higher coronary risk. However, the association of central adiposity with metabolic variables is not the same at all levels of adiposity in all ethnic groups.

This study indicated that there is a significant positive association of WC and WHR with TC, VLDL-C, FPG and FTG among Bengalee men. This implied that among Bengalee men, WC and WHR are associated with an enhanced CHD risk factor profile. Bose and Mascie-Taylor²⁰ in a study conducted in Peterborough, UK, also found that WHR was significantly related to TC among European and migrant Pakistani men. Similar results have been reported from other ethnic groups like African Americans¹¹ and native British.³⁰

Moreover, based on the results of correlation analyses and ANOVA, the present study provided clear evidence that the BMI does not have a significant association with various metabolic risk factors of CHD. Bose and Mascie-Taylor¹⁷ had also reported that the BMI was not significantly associated with TC and FPG among migrant Indians in Britain. It therefore appears that among Indians, both migrants¹⁷ as well as residents in India (present study), central adiposity is more strongly associated with metabolic risk factors of CHD than BMI.

The presence of a significant positive relationship between WHR and CHD risk factors has limited epidemiological applications unless a specific cut-off point of WHR is used. A specific WHR cut-off point allows direct inter-individual as well as inter-population comparisons. In one study, Dasgupta and Hazra²⁴ have utilised the JNC cut-off point of WHR>0.95 to define central obesity among the Bengalee population.

The present study found that CO men had significantly greater means compared with CNO subjects for several metabolic risk factors for CHD. Results of ANOVA using central obesity status (CNO=no, CO=yes) and BMI (BMI tertiles) as categorical variables further demonstrated the significant effect of central obesity, and not of BMI, on these metabolic risk factors. Central obesity status had significant impact on these metabolic variables irre-

spective of the level of BMI. Moreover, there was no significant BMI tertile-central obesity status interaction. These results therefore provide evidence that the JNC cut-off point of WHR>0.95 can be utilised to identify individuals who are at greater risk of developing CHD among Bengalee men.

The only previous study from India²² that compared metabolic risk factor profiles of CHD between CNO and CO subjects did not report any significant difference between the two groups. However, they had used a cut-off point of WHR=0.88 (median WHR) to define CNO and CO individuals; this is not an established cut-off point. Moreover, it may not be appropriate to use WHR=0.88 for use among adult Bengalee men since Gupta and Majumdar's study was not undertaken on Bengalees.²² A further reason for recommending a cut-off point

of WHR>0.95 among Bengalee men is that it would enable the comparison of prevalence rates of central obesity as well as lipid abnormalities associated with it in this population with data available from other ethnic groups.

Since there is vast ethnic heterogeneity in India, future studies should determine whether this cut-off point of 0.95 is 'ethnic-specific' to Bengalees or is applicable to others resident in India. Furthermore, similar studies on Indian women are needed to test the validity of the JNC¹⁴ recommendations for this gender (WHR>0.85). The Indian Diaspora should be studied to determine whether the JNC guidelines can also be applied to them. The Indian Diaspora offers a unique opportunity to study the 'gene-environment' interaction involved in the aetiology of CHD. Although investigations from Britain^{18,20} have studied the relationship of

Table 3**ANOVA of central obesity status (CNO=no, CO=yes) and BMI tertiles with metabolic variables**

Dependent variable	F	P
TC		
BMI tertile	2.072	0.130
Central obesity status	4.270	0.041
BMI tertile-central obesity status interaction	0.007	0.993
VLDL		
BMI tertile	0.180	0.836
Central obesity status	5.488	0.021
BMI tertile-central obesity status interaction	0.186	0.831
FPG		
BMI tertile	0.193	0.825
Central obesity status	5.826	0.017
BMI tertile-central obesity status interaction	0.275	0.760
FTG		
BMI tertile	0.180	0.836
Central obesity status	5.488	0.021
BMI tertile-central obesity status interaction	0.186	0.831

BMI = body mass index

TC = total cholesterol

VLDL-C = very low density lipoprotein cholesterol

FPG = fasting blood glucose

FTG = fasting triglyceride

WHR with CHD risk factors among migrant South Asians, no study has been undertaken on migrant Indians specifically using the JNC guidelines¹⁴ for WHR as a marker for central obesity. Such studies when done in comparison with the native populations should yield valuable information on the ethnic susceptibility to central obesity and its relationship to CHD. Most importantly, prospective studies should be undertaken to obtain the optimum cut-off point to define central obesity among various Indian populations.

ACKNOWLEDGEMENTS

Dr G Hariharan and Dr N Ghosh of Eastern Railways, are gratefully acknowledged for granting permission as well as providing the necessary facilities to undertake this study. Dr Kaushik Bose was a recipient of a Senior Research Associateship of the Council for Scientific and Industrial Research, Government of India. The technical staff of the Department of Pathology, BR Singh Hospital are acknowledged for their help during data collection. Grateful thanks are

also due to Departmental Special Assistance Phase-II, University Grants Commission Programme, Department of Anthropology, Calcutta University, for providing financial assistance. Professor Ranjana Roy, Head of Department, Department of Anthropology, University of Calcutta is acknowledged for providing the necessary facilities to conduct this study.

References

- Kopelman PG. Obesity as a medical problem. *Nature* 2000;404:635-43
- Vague P. The degree of masculine differentiation of obesities: a factor determining predisposition to diabetes, atherosclerosis, gout, and uric-calculus disease. *Am J Clin Nutr* 1956;4:20-34
- Hartz AJ, Rupley DC, Rimm AA. The association of girth measurements with disease in 32,856 women. *Am J Epidemiol* 1984;119:71-80
- Lundgren H, Bengtsson C, Blohme G, Lapidus L, Sjostrum S. Adiposity and adipose tissue distribution in relation to incidence of diabetes in women: results from a prospective population study in Gothenburg, Sweden. *Int J Obes* 1989;13:413-22
- Rimm EB, Stampfer MJ, Giovannucci E, Ascherio A, Spiegelmen D, Colditz GA, et al. Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol* 1995;141:1117-27
- Cox BD, Wichelow MJ, Ashwell M, Prevost AT, Lejeune SR. Association of anthropometric indices with elevated blood pressure in British adults. *Int J Obes* 1997;21:674-80
- Wei M, Gaskill SP, Haffner SM, Stern MP. Waist circumference as a best predictor of NIDDM compared to BMI, WHR and other anthropometric measures in Mexican-Americans: a seven year prospective study. *Obes Res* 1997;5:16-23
- Howard BV, Lee ET, Cowan LD, Fabsitz RR, Howard WJ, Oopik AJ, et al. Coronary heart disease prevalence and its relation to risk factors in American Indians. The Strong Heart Study. *Am J Epidemiol* 1995;142:254-68
- Welty TK, Lee ET, Yeh JL, Cowan LD, Go O, Fabsitz RR, et al. Cardiovascular disease risk factors among American Indians. The Strong Heart Study. *Am J Epidemiol* 1995;142:269-87
- Frisancho AR, Smith S, Albalak R. Relationship of serum cholesterol and truncal body fat distribution among Mexican-Americans is accentuated by obesity. *Am J Hum Biol* 1994;6:51-9
- Kim KS, Robbins D, Turner M, Adams-Campbell LL. Anthropometric determinants of risk factors in an African American population. *Am J Hum Biol* 1998;10:249-58
- Hodge AM, Dowse GK, Erasmus RT, Spark RA, Nathaniel K, Zimmet PZ, et al. Serum lipid and modernization in coastal and highland Papua New Guinea. *Am J Epidemiol* 1996;144:1129-42
- Fiori G, Facchini F, Pettener D, Rimondi A, Battistini N, Bedogni G. Relationships between blood pressure, anthropometric characteristics and blood lipids in high- and low-altitude populations from Central Asia. *Ann Hum Biol* 2002;27:19-28
- The Fifth Report of the Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure (JNC-V). *Arch Intern Med* 1993;153:154-83
- Enas EA, Yusuf S, Mehta JL. Prevalence of coronary artery disease in Asian Indians. *Am J Cardiol* 1992;70:945-9
- Bose K. Coronary heart disease among migrant South Asians in Britain: an overview. *J Ind Anthropol Soc* 1997;32:269-75
- Bose K, Mascie-Taylor CG. Interrelationships of age and the body mass index with risk factors of non-insulin dependent diabetes in European and migrant Asian males. *Am J Hum Biol* 1997;9:291-6
- McKeigue PM, Shah B, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet* 1991;337:382-6
- Bose K, Mascie-Taylor CG. Relationship of body mass index, regional adiposity and body fat distribution with some risk factors of NIDDM in adult Caucasian and migrant Pakistani males. *Bull Mem Soc d'Anthropol Paris* 1997;9:279-90
- Bose K, Mascie-Taylor CG. Conicity index and waist-hip ratio and their relationship with total cholesterol and blood pressure in middle-aged European and migrant Pakistani men. *Ann Hum Biol* 1998;25:11-16
- Gupta R, Mehrishi S. Waist-hip ratio and blood pressure correlation in an urban Indian population. *J Indian Med Ass* 1997;95:412-5
- Gupta R, Majumdar S. Correlation of waist-hip ratio with coronary heart disease and risk factor prevalence in a rural male population. *Indian Heart J* 1994;46:145-8
- Pais P, Pogue J, Gerstein H, Zachariah E, Savitha D, Jayprakash S, et al. Risk factors for acute myocardial infarction in Indians: a case-control study. *Lancet* 1996;348:358-63
- Dasgupta S, Hazra SC. The utility of waist circumference in assessment of obesity. *Indian J Public Health* 1999;43:132-5
- Lohman TG, Roche AF, Martorell R. Anthropometric Standardization Reference Manual. Chicago: Human Kinetics Books, 1988
- Wildman RE, Medeiros DM. Advanced Human Nutrition. Boca Raton, Florida: CRC Press Inc, 2000. p. 332
- Varley H, Gowenlock AH, Bell M. Enzymes. Practical Clinical Biochemistry. Volume 1. Fifth edition. London: William Heinemann Medical Books Ltd, 1980. pp. 625-84
- Friedewald WT, Levy RE, Fredrickson DS. Estimation of the concentration of low density lipoprotein cholesterol in plasma without use of the preparative ultracentrifuge. *Clin Chem* 1972;18:499-502
- Kopelman PG. Causes and consequences of obesity. *Med Int* 1994;22:385-8
- Yasmin, Mascie-Taylor CG. Adiposity indices and their relationship with some risk factors of coronary heart disease in middle-aged Cambridge men and women. *Ann Hum Biol* 2000;27:239-48

Journal of The Royal Society for the Promotion of Health

Panel of Referees

We would like to express our warm appreciation of the substantial time and expertise that our referees have provided over the last year, helping to enhance the quality of published papers. (Some referees have preferred to remain anonymous.)

Mohammad Ameen

Graham Aston

Madan R Bahl

Steve Barrass

Sandra Barry

David Bender

Jim Biggs

Mabel Blades

Andrew D Blann

Ibrahim I Bolaji

Joanna Bowtell

Harry Brewer

Geoffrey Brundrett

Amanda Button

Martin Chaplin

Neil Charman

Rakesh Chopra

Vassiliki Costarelli

Derek Cruickshank

Derrick Crump

Carole A Cull

G Jill Davies

John W T Dickerson

Christine Dickinson

David Donaldson

Raymond J Donaldson

Damien Downing

David Edwards

Toyin Ejidokun

M Elisaf

David J Finney

Philip Fleming

Len Goldstone

Adrian Gossage

Dinah Gould

John Guillebaud

Mike Halls

Harold D Harvey

Anthony Hawkes

Roger Hoare

Selwyn Hodge

Sue Holmes

Stuart J Horner

David Jeffs

Hugh H John

A Mac Johnston

Thomas E Jordan

William Kelly

David Lane

Jeremy Leach

Sammy Lee

Anthony R Leeds

Alexander K C Leung

Helen J Lightowler

Maria Lis-Balchin

Doug MacInnes

Kevin Maguire

Hardipp Mann

Vincent Marks

Anna Maslin

Robert Mason

Peter J May

Andrée le May

Dimitri Mikhailidis

Jane Morgan

Devi R Nair

Carol Noble

Abayomi A Opaneye

Rod Owen

Chris Packham

Paul Palmer

Theresa Phillips

Norman Platt

Bashir Qureshi

William J Reilly

Christopher H Rena

Bryan Reuben

Nagamanickam Sai Sankar

D Shah

Heema Shukla

Sabine Sonnenberg

Antony Stewart

Jill Stewart

Andrew Taylor

Helen Taylor

Steve Tee

Mohammed Twaij

Gerald Vinten

Alexander R P Walker

Susan Ward

Sarah White

William Whitfield

Anthony S Wierzbicki

Richard Willis

Ivan C J Wood